CrossMark



On the determination of vortex ring vorticity using Lagrangian particles

O. Outrata¹, M. Pavelka¹, J. Hron¹, M. La Mantia^{1,†}, J.I. Polanco² and G. Krstulovic²

¹Faculty of Mathematics and Physics, Charles University, Ke Karlovu 3, 121 16 Prague, Czech Republic
²Université Côte d'Azur, Observatoire de la Côte d'Azur, CNRS, Laboratoire Lagrange, Boulevard de l'Observatoire CS 34229, F-06304 Nice CEDEX 4, France

(Received 18 March 2021; revised 14 June 2021; accepted 15 July 2021)

Particles are a widespread tool for obtaining information from fluid flows. When Eulerian data are unavailable, they may be employed to estimate flow fields or to identify coherent flow structures. Here we numerically examine the possibility of using particles to capture the dynamics of isolated vortex rings propagating in a quiescent fluid. The analysis is performed starting from numerical simulations of the Navier-Stokes and the Hall–Vinen–Bekarevich–Khalatnikov equations, respectively describing the dynamics of a Newtonian fluid and a finite-temperature superfluid. The flow-induced positions and velocities of particles suspended in the fluid are specifically used to compute the Lagrangian pseudovorticity field, a proxy for the Eulerian vorticity field recently employed in the context of superfluid ⁴He experiments. We show that, when calculated from ideal Lagrangian tracers or from particles with low inertia, the pseudovorticity field can be accurately used to estimate the propagation velocity and the growth of isolated vortex rings, although the quantitative reconstruction of the corresponding vorticity fields remains challenging. On the other hand, particles with high inertia tend to preferentially sample specific flow regions, resulting in biased pseudovorticity fields that pollute the estimation of the vortex ring properties. Overall, this work neatly demonstrates that the Lagrangian pseudovorticity is a valuable tool for estimating the strength of macroscopic vortical structures in the absence of Eulerian data, which is, for example, the case for superfluid ⁴He experiments.

Key words: vortex dynamics, particle/fluid flow, quantum fluids

1. Introduction

The accurate and well-resolved measurement in space and time of fluid flows represents one of the main ongoing challenges in experimental fluid dynamics. A number of methods

†Email address for correspondence: lamantia@mbox.troja.mff.cuni.cz

O. Outrata and others

have been developed to capture the evolution of velocity fields and other relevant quantities in laboratory flows. A common approach is to visualize the motion of relatively small particles advected by the flow of interest, from which properties of the latter can be inferred (see e.g. Raffel *et al.* 2018). Various techniques can then be employed to quantitatively process the obtained images, the choice mainly depending on the features to be investigated.

The most popular tools are particle image velocimetry (PIV), which allows one to access instantaneous flows fields, i.e. Eulerian data, and particle tracking velocimetry (PTV), which is a Lagrangian technique, focusing on the temporal dynamics of the visualized particles. In principle, both methods can be used to gather complementary information on the same flow but, in practice, this is not always possible. Leaving aside for a moment the important roles played by particle size and inertia, addressed below, the main limiting factor is the particle number density, discussed, for example, by Nobach & Bodenschatz (2009). Specifically, if the number of particles per unit area is smaller than about 0.01 pixel⁻², significant errors occur in PIV measurements. It then follows that, if one cannot seed the flow of interest with enough particles, PTV is the only choice to get quantitative information from the collected images.

An exemplary case is represented by superfluid ⁴He, which is also known as He II (see e.g. Barenghi, Skrbek & Sreenivasan 2014; Mongiovì, Jou & Sciacca 2018). The cryogenic flows of this unique liquid are routinely visualized and several PTV studies have been reported to date – see Švančara *et al.* (2021) for a recent example. However, at a given time, one can usually identify the positions of about 100 particles within a 1 Mpixel image, each particle having an apparent size of a few pixels. Consequently, the visualization data presently collected in He II do not match the quality of the PIV data obtained in water or air, i.e. the former data are more suitable for PTV than for PIV, at least if one has in mind quantitative studies.

The observed behaviour is related to the available experimental techniques, which do not allow one to seed the flow of interest with a relatively large number of particles, at a given time. The density of the liquid is equal to about 145 kg m⁻³ (Donnelly & Barenghi 1998) and it is consequently challenging to employ neutrally buoyant particles, as discussed, for example, by Švančara *et al.* (2018). Indeed, the mass density of the flow-probing particles should match that of the liquid but this is yet to be achieved for visualization experiments in He II. It follows that the particles routinely used – made of solid hydrogen or deuterium – disappear from the field of view after some time, of the order of minutes. Additionally, particles are usually injected into the liquid and, before taking measurements, one must wait for the injection flow to disappear. Then, as already said, one is left with about 100 particles that can be tracked for some time, of the order of seconds (the particle size is usually of the order of micrometres).

To illustrate these facts, we display in figure 1 two images obtained from experimental data discussed by Švančara, Pavelka & La Mantia (2020). Macroscopic vortex rings propagating in superfluid ⁴He were studied using the PTV technique, i.e. solid deuterium particles were dispersed in the liquid and illuminated by a planar laser sheet. The ring-induced motions of the particles were collected by a digital camera, in a plane approximately parallel to the ring propagation direction, that is, in a plane approximately corresponding to one of the ring symmetry planes. The white regions correspond to particles reflecting incoming light and the images were obtained by superimposing hundreds of frames, i.e. the white regions can be viewed as preliminary indications of the tracks followed by the particles (heavier than the fluid in this case). Figure 1(*a*) corresponds to the superposition of movie frames in the laboratory coordinate system,



Figure 1. Visualization of particle trajectories induced by a vortex ring in superfluid ⁴He at 1.50 K, with Reynolds number $Re \approx 10^5$; see Švančara *et al.* (2020) for experimental details. The white regions correspond to particles reflecting incoming light and the images were obtained from averaging the light intensity reflected by the particles over hundreds of movie frames. (*a*) The panel is 25 mm wide and 22 mm high; the nozzle ejecting the fluid parcel is located at the bottom centre of the image and the vortex ring moves upward. (*b*) The panel displays (to scale) the same light intensity values as (*a*) in the coordinate system moving with the vortex ring, where the latter time-dependent position and velocity were computed as discussed by Švančara *et al.* (2020).

whereas, in figure 1(b), the white regions have been centred in the reference system moving with the vortex ring.

We clearly appreciate the emergence of holes, which would not be observed for perfect tracers distributed uniformly through the experimental domain. Indeed, even for relatively small particles, when their action on the flow can be neglected, particle inertia might lead to the decoupling of the particle trajectories from the flow streamlines and, in particular, to preferential concentration (see e.g. Maxey 1987; Balachandar & Eaton 2010).

Additionally, in comparison with typical PIV data, very few particles are seen in figure 1, that is, these data cannot be employed for a robust estimate of Eulerian quantities, such as velocity and vorticity fields. Indeed, in PTV experiments, one does not usually have access to the vorticity field because the latter requires a very large number of particles, vorticity being a small-scale quantity that can present sharp variations in time and space. Nevertheless, having access to the vorticity field – or a proxy thereof – can be of great interest for the description of laboratory flows, e.g. for quantitative comparisons between particle trajectories obtained in different experimental conditions.

This research route was specifically followed by Švančara *et al.* (2020) in their PTV study on the propagation of macroscopic vortex rings in He II. The ring-induced motions of the particles were employed to calculate the Lagrangian pseudovorticity field – a proxy for the Eulerian vorticity field, which can be used to quantitatively measure the strength of the generated vortex rings and which can also provide information on the size, position and velocity of these objects.

Additionally, Švančara *et al.* (2020) have shown that, in conditions yet to be met in experiments, the Lagrangian pseudovorticity θ is equal to half of the the Eulerian vorticity ω . The present work further clarifies the relation between θ and ω , especially in conditions closer to those encountered in typical experiments. We not only consider in the following

how particle concentration affects this relation but also take into account the role of particle inertia.

To this end, we numerically study the motion of particles induced by vortex rings in two different settings. In the first part of the work, we investigate the behaviour of an isolated vortex ring propagating in a Newtonian fluid using the finite element method (FEM) (Brezzi & Fortin 1991) with the incremental pressure correction scheme (Guermond, Minev & Shen 2006). We then compute the Lagrangian pseudovorticity field from the positions and velocities of fluid particles (tracers) advected by the obtained Eulerian flow fields, as a function of experimentally relevant parameters, such as the number of flow-probing particles and their spatial distribution.

Secondly, we investigate the effect of particle inertia on the relation between pseudovorticity and vorticity, in the specific case of a vortex ring propagating in a superfluid. For this, we perform direct numerical simulations of the Hall–Vinen–Bekarevich–Khalatnikov (HVBK) model, which is often employed to describe the large-scale hydrodynamics of He II. Indeed, the dynamics of relatively small spherical particles in flows of superfluid ⁴He has been recently investigated within this theoretical framework (Polanco & Krstulovic 2020).

In summary, the results presented below reinforce the view that the Lagrangian pseudovorticity can be used to quantitatively estimate the strength of relatively large vortical structures in the absence of Eulerian data, which is specifically the case of experiments in He II. We confirm that the magnitude of θ can be appreciably smaller than that of ω , consistent with previous analytical and experimental estimates (Švančara *et al.* 2020). We also show that, in order to get a closer quantitative agreement between the vorticity and pseudovorticity trends, larger particle concentrations should be employed compared with those commonly used in experiments. Furthermore, our results demonstrate that the Lagrangian pseudovorticity can be used to resolve accurately general features of the observed ring dynamics. Such features include the ring time-dependent position and propagation velocity, especially in the case of tracers and particles with sufficiently low inertia. However, we also report that, at present, θ does not seem suitable for capturing the fine details of the vortex ring structure, that is, for reconstructing the corresponding Eulerian vorticity fields.

2. Lagrangian pseudovorticity

The vortices shed at the edges of relatively large objects oscillating in superfluid ⁴He have been recently visualized in planes approximately parallel to the direction of oscillation, by following the flow-induced motions of small solid particles suspended in the liquid (Duda *et al.* 2015; Duda, La Mantia & Skrbek 2017). The strength of these vortices was estimated from the two-dimensional particle positions and velocities using a purpose-made scalar measure, introduced to quantitatively compare vortices obtained under different experimental conditions because, as already noted, it is in general not possible to obtain flow vorticity fields from sparse Lagrangian data.

This scalar quantity was named Lagrangian pseudovorticity θ and, as mentioned above, it was also used by Švančara *et al.* (2020) in their experimental study of macroscopic vortex rings propagating in superfluid ⁴He. It is here defined as

$$\theta(\mathbf{r},t) = \left\langle \frac{\left[(\mathbf{r}_i - \mathbf{r}) \times \mathbf{u}_i \right]_z}{|\mathbf{r}_i - \mathbf{r}|^2} \right\rangle_{\mathcal{M}},\tag{2.1}$$

where r_i and u_i denote the particle positions and velocities in the considered plane, respectively. The angle brackets indicate the ensemble average within the set \mathcal{M} of

Lagrangian particles, which are captured within a time window centred at time t and are found within an annular region centred, on a chosen grid, at the inspection point r. The subscript z denotes the axis perpendicular to the observation plane, i.e. we consider here only the non-zero component of the vector product (the observed particle tracks occur in a plane). The size of the annular region was chosen in order to have enough particles for the calculation of θ and, at the same time, to exclude diverging contributions from particles too close to the inspection point r.

Before proceeding, it is now useful to note a few properties of the Lagrangian pseudovorticity θ . As mentioned above, it can be shown analytically that θ is closely related to the Eulerian vorticity, in the case of fluid particles that are homogeneously distributed in space (Švančara *et al.* 2020). Furthermore, it was reported that, for a Rankine vortex, the magnitude of θ decreases when the area of the annular region used for its calculation increases, that is, when the outer radius of the region increases, for given grid and number of fluid particles within the annular region (Švančara *et al.* 2020). Additionally, this magnitude can be appreciably smaller than that of ω when the area is significantly larger than the area where the flow vorticity is concentrated. For example, when the region outer radius is 10 (100) times larger than the Rankine vortex radius, the maximum pseudovorticity magnitude, located at the vortex centre, is more than 5 (50) times smaller than the corresponding vorticity value. We note in passing that changing the grid or the number of points within the annular region has a less significant effect on the calculated pseudovorticity magnitude in comparison with the just mentioned influence of the annular region size.

A similar outcome – displayed in figure 2 – can be obtained for a Lamb–Oseen vortex, which is characterized by a vorticity spatial distribution closer to that observed for vortex rings (see, for example, figure 4 in § 3.2, where relevant numerical results are plotted). It is also apparent that the pseudovorticity may change sign in the vortex vicinity, similarly to what is seen in figure 4(b). Additionally, it is evident from the inset of figure 2 that the number of fluid particles distributed in the annular region becomes important solely when it assumes relatively small values. The same applies to the chosen grid size that affects the computed pseudovorticity values only for numbers of grid points smaller than those considered in the figure.

Before presenting our numerical results, we also note here that in Appendix A, additional analytical results on the relation between pseudovorticity and vorticity are reported. Specifically, we show that, for an isolated vortex having a purely azimuthal velocity field, the pseudovorticity magnitude becomes smaller if the area employed for its estimate increases. We also suggest an explanation for the sign changes of the pseudovorticity displayed in figures 2 and 4, that is, the outcome can be explicitly related to the fact that the corresponding particle distributions are not homogeneous or isotropic.

3. Finite element method ring

The finite element method (FEM) (Brezzi & Fortin 1991) was chosen for the space discretization of the well-known equations governing the isothermal dynamics of an incompressible viscous fluid. We specifically employed the Taylor–Hood mixed FEM using the FEniCS library (Alnæs *et al.* 2015). The discretization in time was performed employing the implicit Crank–Nicolson scheme and the incremental pressure correction scheme (Guermond *et al.* 2006) was used for finding an efficient approximate solution of the coupled velocity–pressure problem.



Figure 2. Pseudovorticity for a Lamb–Oseen vortex as a function of the distance from the vortex axis (radius). Vortex parameters: circulation $\Gamma = 50 \text{ mm}^2 \text{ s}^{-1}$, kinematic viscosity $\nu = 0.01 \text{ mm}^2 \text{ s}^{-1}$ and time t = 10 s. Black circles: vortex velocity ν ; red circles: vorticity ω ; open circles: pseudovorticity θ . The corresponding subscripts in the legend indicate the radius, in mm, of the area used for the calculation of θ from (2.1). One thousand equally spaced grid points were employed for the estimate shown in the main panel, up to a radius of 10 mm, and 100 fluid particles were distributed in the annular region, equally spaced along the radial direction (the pseudovorticity is not computed in the region centre and is set to zero at the vortex centre). Note that the chosen particle distribution is not radially symmetric, that is, the particles are clustered in the vortex centre vicinity. The results plotted in the inset were obtained using 100 equally spaced grid points, up to a radius of 10 mm, and the number of equally spaced fluid particles within the annular region is specified in the legend. The region outer radius for the data in the inset is 30 mm. The magenta line indicates the pseudovorticity trend obtained using 1000 equally spaced grid points, corresponding to the cyan open circles in the main panel.

A schematic view of the numerical domain employed for the simulations in two dimensions is shown in figure 3. For these simulations we have specifically chosen boundary conditions close to experimental ones (Švančara *et al.* 2020). On all domain boundaries the fluid velocity is set to zero, excluding the top of the domain, where a traction free boundary condition is imposed, and the nozzle top, where the vertical fluid velocity has a spatial parabolic profile during the first second of the simulation (it is zero at later times). There, the fluid peak velocity linearly increases from 0 to 10 mm s⁻¹ during the first 0.1 s, remains constant for the following 0.8 s and linearly decreases from 10 to 0 mm s⁻¹ during the last 0.1 s of the first second of the simulation.

The latter boundary condition simulates the vortex ring generation process, which may occur when a parcel of fluid is ejected into a reservoir through a nozzle. In practice, this is often achieved using a piston moving inside the nozzle with a velocity U_p , for a stroke length L. An adequate Reynolds number can then be defined as

$$Re = \frac{U_p L}{2\nu} = \frac{\Gamma_0}{\nu},\tag{3.1}$$

where, for the present simulations, the fluid kinematic viscosity is $\nu = 0.01 \text{ mm}^2 \text{ s}^{-1}$ and $\Gamma_0 = U_p L/2$ indicates the ring nominal circulation (see e.g. Švančara *et al.* 2020).

Here, we set U_p equal to the mean value of the imposed fluid velocity during the first second of the simulation, i.e. $U_p = 6 \text{ mm s}^{-1}$, and the stroke length $L = U_p t_p = 6 \text{ mm}$,



Figure 3. Two-dimensional domain used for the FEM simulations, composed of the grey, light blue and red areas. The yellow area -5 mm wide and 10 mm high – represents the nozzle and is not included in the numerical domain (a parcel of fluid is ejected through the nozzle to generate the vortex ring). The grey area is 50 mm wide and 110 mm high. In the light blue area -20 mm wide and 79 mm high -5676 fluid particles are uniformly distributed at the beginning of the simulation (experiment particle distribution; the area is located 1 mm above the nozzle and includes the red area). In the red area -10 mm wide and 5 mm high $-22\,801$ fluid particles are uniformly distributed 1 s after the beginning of the simulation (delay particle distribution; the area is located 1 mm above the nozzle). The origin of the Cartesian reference system (indicated by black arrows) is at the bottom centre of the yellow area.

where $t_p = 1$ s is the imposed duration of the fluid ejection into the reservoir. It follows that $\Gamma_0 = 18 \text{ mm}^2 \text{ s}^{-1}$ and Re = 1800, which means that our FEM ring is most likely in the laminar regime (Glezer 1988). Additionally, L/D = 1.2, where D = 5 mm is the nozzle internal diameter, that is, we do not expected a prominent ring wake (Gharib, Rambod & Shariff 1998); note also that the ring nominal impulse, per unit of density, $I_0 = \Gamma_0 \pi (D/2)^2 = 353 \text{ mm}^4 \text{ s}^{-1}$ (Sullivan *et al.* 2008).

A mesh independence check was performed on three meshes with different densities for the space discretization, resulting in systems with sizes ranging from 2×10^5 to 1.3×10^6 degrees of freedom. The size of the time step ranged from 0.01 to 0.001 s. Full three-dimensional computations were also performed for the sake of comparison and the obtained results were not appreciably different from the two-dimensional ones; see Outrata (2020) for further details. In the following, we discuss the two-dimensional numerical results obtained using the most refined mesh, that with approximately 1.3×10^6 degrees of freedom, and with the smallest time step, equal to 0.001 s (the entire simulation lasts 15 s).

3.1. Initial fluid particle distributions

We considered fluid particles in the obtained Eulerian flow fields and tracked their motions in order to compute – from the particle positions and velocities – the Lagrangian pseudovorticity, (2.1). As a first step, we uniformly distributed, at the beginning of the simulation, approximately 6000 particles in a relatively large area of the numerical domain, shown as the light blue region in figure 3, our aim being to replicate actual experimental conditions (Švančara *et al.* 2020). We refer to this arrangement as the experiment particle distribution and label it with the letter E. As shown in the following, its results are characterized by the fact that many particles do not move, being away from the vortex ring, similarly to what is observed in experiments.

O. Outrata and others



Figure 4. Pseudovorticity and vorticity maps obtained 10 s after the beginning of the FEM simulation. Pseudovorticity computed from (a) the experiment (E) and (b) the delay (D) particle distributions. (c) Vorticity field. The reference system is that defined in figure 3. Clockwise fluid rotation corresponds to positive values of pseudovorticity and vorticity (the ring is moving upward). Note that the vorticity and pseudovorticity values in the nozzle proximity, for vertical positions smaller than 15 mm, are not taken into account in the other calculations reported here.

For our analysis we also employed another particle arrangement and refer to it as the delay distribution, because the fluid particles were added to the Eulerian flow fields 1 s after the beginning of the simulation (we label this distribution with the letter D). Additionally, compared with E, more particles were distributed in a smaller area, shown as the red region in figure 3, our main aim being to increase the number of particles significantly contributing to the pseudovorticity computation, that is, the number of moving particles.

The inner and outer radii of the annular region employed for pseudovorticity estimates, (2.1), were set to 0.1 and 3 mm, respectively, and the corresponding time window is 10 ms wide. For both distributions, the chosen grid is a rectangular mesh of 53×126 inspection points, covering a 20.8 mm wide and 60.0 mm high area of the computational domain, centred just above the nozzle (see figure 3). Note that on the same grid the obtained flow vorticity was interpolated and that the fluid particle velocities are known from the numerical simulation, that is, they are not computed from the particle positions as in the experiments (Švančara *et al.* 2020).

3.2. Pseudovorticity maps

Having set these conditions, we calculated the Lagrangian pseudovorticity θ from the instantaneous positions and velocities of both particle distributions. In figure 4(a,b) we display the θ maps obtained 10 s after the beginning of the simulation. In figure 4(c) the same-time flow vorticity is plotted.

If one compares figures 4(a) and 4(c), it is apparent that the vortex ring vertical position obtained from the experiment particle distribution is quite close to that resulting from the vorticity map. On the other hand, the ring spatial extent for the E distribution is appreciably larger than that apparent from the vorticity map. Additionally, the experiment pseudovorticity magnitudes are at least 100 times smaller than the vorticity ones.

Instead, for the delay particle distribution the magnitudes of θ are significantly closer to those of ω (see figure 4*b*,*c*). More importantly, the corresponding spatial



Figure 5. Root-mean-square (r.m.s.) pseudovorticity (red) and vorticity (black) profiles obtained 10 s after the beginning of the FEM simulation. The pseudovorticity is computed from the delay (D) particle distribution. Profiles are averaged either in the vertical (*a*) or horizontal (*b*) directions. The different scales on the vertical axes are for pseudovorticity (left axes) and vorticity (right axes). The reference system is as in figure 4. Note that the vorticity and pseudovorticity values in the nozzle proximity, for vertical positions smaller than 15 mm, are not taken into account in the other calculations reported here.

distributions are quite similar to each other. This is further confirmed in figure 5, which shows root-mean-square profiles of θ and ω , respectively averaged over the vertical and horizontal directions.

We note in passing that particle positions characterized by velocities smaller than 0.01 mm s^{-1} were not taken into account for the present pseudovorticity estimate because this led to (slightly) smoother θ maps and (slightly) larger pseudovorticity magnitudes for the E particle distribution. The effects were, however, absent for the D particle distribution, due to the larger velocities of the considered particles. Additionally, for the D distribution, the vorticity in the nozzle proximity is not seen, because the particles are distributed 1 s after the beginning of the simulation.

More generally, the results displayed in figures 4 and 5 confirm that the Lagrangian pseudovorticity can be used to characterize vortex ring features in the absence of Eulerian data (Švančara *et al.* 2020). Additionally, they demonstrate that the quantitative agreement between θ and ω strongly depends not only on the number of flow-probing particles, but also on how they are distributed in the region of interest.

3.3. Ring position, velocity and radius

In order to further support the previous claims, we calculated other relevant quantities from the obtained pseudovorticity and vorticity maps, which enable, for example, the estimation of the vortex ring location and size over time.

Dataset	$\begin{pmatrix} \theta_0 \\ (s^{-1}) \end{pmatrix}$	v (mm s ⁻¹)	v/U_p	2 <i>R</i> (mm)	$\frac{C}{(\mathrm{mm}^2 \mathrm{s}^{-1})}$	A_C (mm ²)
Vorticity	0.800	2.08 ± 0.13	0.35	3.09 ± 0.14	46.21 ± 2.32	13.78 ± 0.61
Tracer D	0.070	2.09 ± 0.24	0.35	3.37 ± 0.09	5.37 ± 0.61	16.95 ± 2.58
Tracer E	0.002	2.06 ± 0.42	0.34	4.51 ± 0.23	0.28 ± 0.04	35.06 ± 1.91

Table 1. Properties of the FEM ring. (i) Chosen threshold θ_0 for ring identification, set to approximately 10% of the maximum magnitude at late times. (ii) Ring vertical velocity v; see also figure 6(b). (iii) Ratio between the mean value of the ring vertical velocity v and the piston velocity U_p , which is found to be consistent with values reported in the literature (see e.g. Sullivan *et al.* 2008). (iv) Ring diameter 2*R*; see also figure 7. (v) Ring circulation *C* from (3.2); see also figure 8. (vi) Ring area A_C ; see also figure 9. Results obtained by applying the chosen threshold values. Symbols as in Švančara *et al.* (2020).



Figure 6. Ring position (*a*) and velocity (*b*) as a function of the time *t* elapsed from the beginning of the FEM simulation. The reference system is that defined in figure 3.

As a first step, following Švančara *et al.* (2020), we chose, for each map, a positive threshold value θ_0 that is employed to identify and track the vortex ring. The positive (clockwise) part of the ring is then made of grid points with $\theta > \theta_0$; the negative (counterclockwise) one is instead described by $\theta < -\theta_0$. The chosen values – listed in table 1 – are constant in time and are set to approximately 10 % of the maximum magnitude at late times.

Once the filtered maps are computed, we can identify the centres of the positive and negative vortical regions, and the distance between them, which is known as the ring diameter 2R (Gan & Nickels 2010). Additionally, the ring position P is defined as the centre of the two vortices and its velocity is calculated by convolving P with a suitable Gaussian kernel (Švančara *et al.* 2020).

We note in passing that the values listed in table 1 are obtained for times ranging between 3.00 and 14.79 s (the entire simulation lasts 15 s). The upper limit is due to the Gaussian algorithm chosen for the ring propagation velocity calculation, while the lower one is



Figure 7. Ring radius R as a function of the time t elapsed from the beginning of the FEM simulation. The reference system is as in figure 6.

imposed to remove the influence on the results of the fluid vorticity in the nozzle proximity (see figure 4) – in the present work we decided not to investigate the ring generation process mainly because this was not accessed in relevant experiments (Švančara *et al.* 2020).

In figure 6 we plot the position and velocity of the vortex ring as a function of the time elapsed from the beginning of the simulation. It is evident that the ring position is well captured by both pseudovorticity maps, while the ring velocity obtained from the experiment particle distribution is characterized by a significantly larger standard deviation compared with that derived from the delay distribution (see also table 1). On the other hand, the corresponding mean values are quite close to each other (see again table 1). Note also that, at sufficiently late times, the ring velocity shows a tendency to decrease in magnitude, which is a behaviour occurring to vortex rings propagating in a viscous fluid (see e.g. Maxworthy 1974; Sullivan *et al.* 2008).

One also expects that the decrease of the ring velocity magnitude is accompanied by an increase of its radius. As shown in figure 7, which displays the vortex radius as a function of the time elapsed from the beginning of the simulation, this behaviour is indeed captured by the delay particle distribution. However, this is not the case for the experiment distribution. The outcome can be related to the above mentioned fact that the ring spatial extent is not well resolved by this particle arrangement (see again figure 4).

We can therefore say that, using the Lagrangian pseudovorticity, one can track rather well the position of a vortex ring and estimate with some confidence its propagation velocity. On the other hand, in order to get quantitative information on the ring spatial extent, one should also employ for the Lagrangian analysis a number of fluid particles significantly larger than those used to date in experiments (Švančara *et al.* 2020). Additionally, these particles should be distributed in a relatively small area, with dimensions comparable with those of the ring itself.



Figure 8. Ring circulation as a function of the time *t* elapsed from the beginning of the FEM simulation. (*a*) The solid lines indicate the ring circulation Γ obtained using the full maps, from (3.2) with $\theta_0 = 0$ (see figure 4). The dashed lines denote the circulation *C* calculated using the filtered maps, from (3.2) with the threshold values listed in table 1. (*b*) Circulation *C* normalized by the ring nominal circulation Γ_0 . Note that in (*b*) and in all the other FEM figures – excluding figures 4 and 5 – the chosen pseudovorticity and vorticity thresholds are taken into account for the estimate of the plotted quantities.

3.4. Ring circulation and area

Following Švančara *et al.* (2020) we also estimate the ring circulation C from the pseudovorticity (and vorticity) maps. For this, we use the relation

$$C = \frac{C_{+} - C_{-}}{2} = \frac{\sum(\theta_{+}a) - \sum(\theta_{-}a)}{2},$$
(3.2)

where the summations only include the grid points characterized by values of θ (or ω) passing the threshold θ_0 listed in table 1. The subscript + (-) denotes that the pseudovorticity and vorticity values are positive (negative), and *a* is the fixed area associated with each grid point. Since $C_+ \approx |C_-|$, we discuss here only their mean absolute value *C*.

In figure 8(a) the dashed lines represent the temporal evolution of *C* and the solid lines correspond to the actual ring circulation Γ , obtained using the full maps, that is, with the threshold θ_0 set to zero. We can see that *C* is consistently smaller than Γ , likely because the filtered maps do not capture the wake shed by the moving ring. Figure 8(a) also displays the tendency of the ring circulation to decrease at late times. This is an expected behaviour for both turbulent and laminar rings, as discussed, for example, by Maxworthy (1974) and Didden (1979), respectively.

Additionally, the circulation magnitude derived from the flow vorticity (black line) is approximately 100 times larger than that obtained from the experiment particle distribution (blue line). This is also evident from figure 8(*b*), which displays *C* normalized by the ring nominal circulation Γ_0 . A similar outcome was reported by Švančara *et al.* (2020), who



Figure 9. Ring area as a function of the time t elapsed from the beginning of the FEM simulation.

showed that experimentally obtained values of *C* can be much smaller than relevant values of Γ_0 , up to approximately two orders of magnitude. Note that for these experiments it was not possible to access the actual ring circulation Γ and that the nominal circulation was estimated from suitable values of fluid velocity.

From figure 8(b) it is also apparent that $C/\Gamma_0 \approx 2$, when C is computed from the filtered vorticity maps (black line). Indeed, measured values of ring circulation were found to be of the same order as Γ_0 by a few investigators, in various experimental conditions (see e.g. Maxworthy 1974; Didden 1979; Borner & Schmidt 1985). Note, however, that the ring nominal circulation $\Gamma_0 = U_p^2 t_p/2$; that is, if we set, for our FEM ring, $U_p = 10 \text{ mm s}^{-1}$ and $t_p = 0.8 \text{ s}$, we obtain a nominal circulation more than two times larger than that employed in figure 8(b) – see also the above discussion of (3.1). It then follows that the numerical values shown in the figure should solely be regarded as first-order estimates, considering also that, as discussed, for example, by Glezer (1988), the nozzle geometry and the piston velocity time history may have a significant influence on the resulting ring features.

Following Švančara *et al.* (2020) we can now estimate the ring area A_C from (3.2), by setting $\theta_+ = 1$ and $\theta_- = -1$. We plot the result in figure 9(*a*) as a function of the time elapsed from the beginning of the simulation. Additionally, in figure 9(*b*), we display the ratio A_C/A_R , where $A_R = \pi R^2$ and *R* is the estimated ring radius shown in figure 7.

The most prominent feature of figure 9 is likely the peak observed for the delay particle distribution (red lines) at short times. This can be related to the fact that, at around the same time, the ring wake disappears from the filtered θ maps. Note that a similar peak is also observed in the estimated vertical velocity of the ring, at $t \approx 6$ s (figure 6b).

Additionally, from figure 9(b), it appears that the ring radius R can be used as a valuable first-order estimate of the ring spatial extent. More importantly, we see from both panels that the ring spatial extent is quite well captured by the delay particle distribution, consistent with our previous discussion.

In summary, our results confirm that the Lagrangian pseudovorticity can be used to quantify general features of relatively large vortical structures – such as their position

and propagation velocity – and that this is especially meaningful when Eulerian data are not available, which is specifically the case for experiments in superfluid ⁴He (Švančara *et al.* 2020). Additionally, we have shown here that the spatial extent of these vortices can be quantitatively determined by θ maps solely in conditions yet to be met in relevant experiments, that is, only for sufficiently dense particle distributions, which, by the way, could allow one to directly access Eulerian flow fields.

So far we have considered fluid particles, which by definition exactly follow the flow, but the particles commonly used in experiments can deviate from streamlines due to their inertia. In the next section, we then investigate the effect of particle inertia on the relation between θ and ω .

4. Hall-Vinen-Bekarevich-Khalatnikov ring

The Hall–Vinen–Bekarevich–Khalatnikov (HVBK) model is often used to describe the large-scale hydrodynamics of superfluid ⁴He. At finite temperatures, between approximately 1 and 2.2 K, this unique liquid can be seen as a mixture of two components: the superfluid component, with no viscosity, and the normal fluid component, with Newtonian viscosity (see e.g. Landau 1941; Donnelly 2009). The HVBK model is valid at scales much larger than the typical distance between quantized vortices, which are line singularities – holes – within the superfluid component. The model therefore describes the coarse-grained dynamics of the two components. Within this framework, their respective velocity fields v_s and v_n follow two coupled incompressible Navier–Stokes equations:

$$\frac{\partial \boldsymbol{v}_n}{\partial t} + (\boldsymbol{v}_n \cdot \nabla) \boldsymbol{v}_n = -\frac{\nabla p_n}{\rho_n} + \nu_n \nabla^2 \boldsymbol{v}_n - \frac{\rho_s}{\rho_n} \boldsymbol{f}_{ns}, \qquad (4.1)$$

$$\frac{\partial \boldsymbol{v}_s}{\partial t} + (\boldsymbol{v}_s \cdot \nabla) \boldsymbol{v}_s = -\frac{\nabla p_s}{\rho_s} + \nu_s \nabla^2 \boldsymbol{v}_s + \boldsymbol{f}_{ns}, \qquad (4.2)$$

$$\nabla \cdot \boldsymbol{v}_s = \nabla \cdot \boldsymbol{v}_n = 0. \tag{4.3}$$

Here, ρ_s and ρ_n are the temperature-dependent densities of the two fluids (the total fluid density is $\rho = \rho_n + \rho_s$), while p_s and p_n indicate their respective pressure fields.

The two fluids are coupled by a mutual friction force f_{ns} . This force is responsible for the transfer of energy between the components, and it also contributes to the dissipation of energy of the entire system. As a first approximation, f_{ns} increases linearly with the relative velocity between the components, that is, $f_{ns} = \alpha \Omega_0 (v_n - v_s)$, where α is a non-dimensional mutual friction coefficient - tabulated by Donnelly & Barenghi (1998) and Ω_0 indicates a mutual friction frequency, related to the density and orientation of the quantized vortices embedded in the superfluid. For example, in isotropic superfluid turbulence, this frequency may be taken as proportional to a characteristic superfluid vorticity (L'vov, Nazarenko & Skrbek 2006). Here, as we consider the case of an isolated vortex ring, this frequency is taken as a constant control parameter. We have also included in (4.2) a superfluid dissipation term, parameterized by an effective temperature-dependent kinematic viscosity v_s (Vinen & Niemela 2002). This term accounts for the dissipation of superfluid energy at scales smaller than those resolved by the HVBK model. Such dissipation mechanisms include quantized vortex reconnections and excitation of Kelvin waves, which transport energy towards the smallest scales of the system (see e.g. Švančara & La Mantia 2019; Villois, Proment & Krstulovic 2020). Additionally, in (4.1) the kinematic viscosity of the normal component ν_n is set equal to μ_n/ρ_n , where μ_n is the fluid dynamic viscosity, also tabulated by Donnelly & Barenghi (1998).

924 A44-14

In the reference study by Švančara *et al.* (2020), solid deuterium particles were used to evaluate the Lagrangian pseudovorticity associated with vortex rings generated in superfluid ⁴He. Such particles are not perfect tracers of the flow, as their density is about 1.4 times larger than that of the fluid, and their typical diameter is a few micrometres (Švančara & La Mantia 2017). Hence, particle inertia is expected to play some role in pseudovorticity estimates.

In superfluid ⁴He, inertial particles are not only subject to the Stokes drag associated with the normal fluid viscosity, but also respond to pressure gradient forces from the two fluids (see e.g. Sergeev & Barenghi 2009). Finite-size effects can be neglected, as particles are much smaller than the core size of the studied vortex ring, as reported below, in § 4.1. Consequently, if one also neglects Basset history terms and assumes that particles are spherical, the particle equation of motion reads

$$\frac{\mathrm{d}\boldsymbol{v}_p}{\mathrm{d}t} = \frac{\boldsymbol{v}_n(\boldsymbol{x}_p) - \boldsymbol{v}_p}{\tau_p} + \beta \left(\frac{\rho_n}{\rho} \frac{\mathrm{D}\boldsymbol{v}_n}{\mathrm{D}t} + \frac{\rho_s}{\rho} \frac{\mathrm{D}\boldsymbol{v}_s}{\mathrm{D}t}\right),\tag{4.4}$$

where x_p and v_p are the particle position and velocity, respectively, and D/Dt denotes the material derivative associated with each fluid component. The density parameter β accounts for added mass effects and is given by $\beta = 3\rho/(2\rho_p + \rho)$, where ρ_p is the particle density. The Stokes time, representing the typical particle response time to normal fluid fluctuations, is $\tau_p = a_p^2/(3\beta v)$, where a_p denotes the particle radius and $v = (\rho_n/\rho)v_n =$ μ_n/ρ indicates the temperature-dependent kinematic viscosity of superfluid ⁴He, also tabulated by Donnelly & Barenghi (1998). Formally, tracer dynamics is obtained in the limit $\tau_p \rightarrow 0$.

The HVBK equations (4.1)-(4.3) are solved by direct numerical simulation in a three-dimensional periodic domain using a Fourier pseudo-spectral solver (Homann *et al.* 2009). Particles are initialized at random locations in the domain and are advanced in time according to (4.4). Fluid velocities and accelerations are interpolated at particle positions using fourth-order B-splines (van Hinsberg *et al.* 2012). Time advancement of fields and particles is performed using a third-order Runge–Kutta scheme.

Figure 10 displays results from a typical simulation, in which we observe the vortex ring together with spherical particles having the same density of deuterium particles, at the chosen temperature T = 1.9 K (see numerical details later). All panels correspond to the same time of ring evolution, for different particle families. From figures 10(a) to 10(c) the particle size is increased, leading to a larger Stokes response time. As a consequence, the effect of inertia becomes apparent and holes (regions without particles) are clearly observed. Such structures are similar to those seen in the experimental pictures displayed in figure 1, but one should keep in mind that the latter are not Eulerian images, that is, they were not obtained at a fixed time as the images in figure 10.

4.1. Simulation set-up

We consider superfluid ⁴He at 1.9 K. At this temperature, the densities of the normal fluid and superfluid components satisfy the ratio $\rho_n/\rho_s = 0.72$ and the liquid kinematic viscosity is $\nu \approx 0.01 \text{ mm}^2 \text{ s}^{-1}$. This value is very close to that used for the FEM ring – note that, as stated above, the density and viscosity of He II depend on temperature (see e.g. Donnelly & Barenghi 1998). Additionally, at this temperature, the effective superfluid viscosity is close to that of the normal fluid (Vinen & Niemela 2002) and the mutual friction coefficient is $\alpha = 0.206$. For the numerical simulations we set the mutual friction



Figure 10. HVBK simulation of a vortex ring moving downward with three families of deuterium particles at a relatively early time, approximately 1 s after the beginning of the simulation. (*a*) Particles with radius of 5 μ m. (*b*) Particles with radius of 10 μ m. (*c*) Particles with radius of 30 μ m. The respective Stokes numbers are $St \approx 0.1$, 0.3 and 2.5 (see table 2). The peculiar shape observed in (*c*) is due to particles being expelled from highly rotating regions as a result of their high inertia. Additionally, particles initially located in front of the vortex ring are propelled upwards via the ring centre in a ballistic motion, resulting in the observed wake pattern.

parameter to $\Omega_0 \approx 110 \text{ s}^{-1}$. We can thus consider that the fluid components are well locked to each other for the time scales relevant to this work.

An isolated vortex ring of normal fluid is introduced in the middle of the periodic box, which has 28 mm sides. The origin of the chosen reference system is located at the box centre. The ring radius is R = 3.5 mm and its initial circulation is $\Gamma_0 \approx 95$ mm² s⁻¹. It follows from (3.1) that the ring Reynolds number is slightly larger than 10⁴. Note that the superfluid component is initially at rest. However, due to mutual friction, the normal fluid vortex ring quickly induces the formation of an equivalent superfluid ring. Such a macroscopic superfluid vortex is typically composed of a bundle of quantized vortices (see e.g. Wacks, Baggaley & Barenghi 2014) that are modelled in the HVBK framework as a continuous vorticity field. In this work we are not interested in this fast transient dynamics.

Note also that, due to the domain periodicity, a vortex ring eventually returns to its initial position. Although the ring leaves a wake behind it, its dynamics is almost unaffected by it, since the wake is rapidly dissipated by viscosity (within one period, lasting about 5 s). However, as particles are not continuously injected in the domain, their spatial distribution becomes inhomogeneous after the passing of a ring. Hence, the studied configuration mimics an experiment in which rings are injected periodically, but one should also account for the fact that the ring size and velocity change with time. We will not further exploit this analogy in the present work.

The vortex ring core is initialized with a Gaussian vorticity profile, $\omega(r) = \omega_0 \exp[-r^2/(2r_0^2)]$, where r is the distance from the core centre. The vortex core radius $r_0 = 0.2$ mm and the vorticity at the core centre is set to fix the value of the ring circulation. We note in passing that the generation of such a thin vortex ring does not appear straightforward with the apparatus employed by Švančara *et al.* (2020), mainly because the experimentally generated rings seem to have a rather large core size, but, as shown below, the HVBK ring provides a good setting to study the effect of inertia.

As detailed in 4.2, we track solid hydrogen and deuterium particles – by setting their density – with sizes similar to those found in ⁴He experiments. For comparison, we

also follow perfect tracers of the normal fluid. We specifically consider three families of spherical particles, having radii of 5, 10 and 30 µm, and we seed 5×10^5 particles, homogeneously distributed over the whole (three-dimensional) computational domain. The corresponding Stokes times range approximately from 1 to 41 ms. The characteristic ring time τ_r is estimated using the initial circulation Γ_0 and the core radius r_0 . We obtain that $\tau_r \approx 17$ ms, i.e. relevant values of Stokes number $St = \tau_p/\tau_r$ are between 0.04 and 2.47 – see also table 2.

4.2. Pseudovorticity maps

At the beginning of the simulation, particles are suspended in the cubic volume. A layer of the latter, parallel to the ring propagation direction and situated in the centre of the periodic box, is considered for the pseudovorticity estimate. This volume layer is 0.2 mm thick and contains between 3100 and 3600 particles at each time step, which lasts approximately 0.09 s (the entire simulation lasts approximately 27 s). The positions and velocities of these particles were then employed for the calculation of the pseudovorticity according to (2.1). This estimation was performed on a square grid of 101×101 inspection points, covering a $27.6 \times 27.6 \text{ mm}^2$ area. The chosen annular region has 0.1 mm inner and 3 mm outer radii, which are the same values used for our FEM analysis in a viscous fluid, discussed in § 3. The corresponding time window is equal to the time step, that is, for the HVBK ring, no temporal averaging was carried out.

Note also that, as in the FEM study, the numerically obtained flow vorticity values were interpolated on the chosen pseudovorticity grid, for the sake of comparison. Additionally, in order to follow the ring motion for distances larger than the box size, the ring is at each time step centred in the box middle using the position of the normal fluid vorticity maximum at that time step (the introduced ring is symmetric). From the latter maximum position it is then possible to get the ring position as a function of time, in a chosen reference system not moving with the ring, as for the FEM ring.

Pseudovorticity maps were obtained for fluid tracers and inertial particles. The latter are characterized by a density different from that of the fluid and by a finite Stokes response time (a finite radius). Additionally, we chose particles having features compatible with recent visualization experiments in superfluid ⁴He (see e.g. Švančara *et al.* 2018). Specifically, we chose spherical particles made of solid deuterium D_2 – heavier than liquid He II, with a density ratio of 1.37 at 1.9 K – and solid hydrogen H₂ – lighter than superfluid ⁴He, with a density ratio of 0.60 at the same temperature. Similarly, we selected particle radii of the same order as and larger than the dimensions of particles in experiments – see La Mantia & Skrbek (2014) and Švančara & La Mantia (2017) for typical distributions of particle sizes.

Pseudovorticity maps calculated from the different particle classes are shown in figure 11. As in the FEM analysis, it is possible to capture the instantaneous ring location using tracer particles, but the magnitude of the computed pseudovorticity is two orders of magnitude smaller than that of the vorticity; see also the corresponding circulation values in table 2. Moreover, the pseudovorticity maps obtained using small inertial particles are quantitatively similar to those obtained using tracers. This is expected, as the motion of small particles should not be strongly affected by inertial effects. More practically, it indicates that hydrogen and deuterium particles having radii up to 10 μ m – corresponding to St < 0.3 – may be used to accurately detect the presence of a vortex ring in the present setting. In our study, it is only for the largest particles, with radius $a_p = 30 \ \mu$ m, corresponding to St > 1, that pseudovorticity estimates importantly deviate from those of



Figure 11. Vorticity and pseudovorticity maps obtained 2.69 s after the beginning of the HVBK simulation. (*a*) The Eulerian vorticity map. (b–h) Pseudovorticity maps obtained using different types of particles, as indicated in the panel titles (the shown dimension indicates the particle radius). The origin of the reference system, moving with the ring, is located at the centre of the periodic box. Clockwise fluid rotation corresponds to positive values of pseudovorticity and vorticity (the ring is moving downward).

tracer particles, even though relevant qualitative information may still be extracted from the resulting maps.

In our simulations, we also observe that, for fluid tracers, the number of particles does not influence appreciably the obtained results. For example, if the chosen layer of the cubic volume is five times thicker, the number of particles used for the pseudovorticity calculation also increases by approximately five times, but the corresponding magnitude does not change appreciably, although its behaviour in space and time is possibly smoother than that obtained for the thinner layer. Note that similar results were obtained for the FEM ring, that is, increasing the number of particles in a given area did not significantly improve the agreement between vorticity and pseudovorticity magnitudes above a certain number of particles (see also the inset of figure 2). We expect that, for given number of particles, a better agreement could be obtained for smaller annuli, as discussed in Appendix A. On the other hand, if one decreases the size of the region used for the pseudovorticity estimate in actual experiments, the number of particles in that region also decreases



Figure 12. Root-mean-square (r.m.s.) pseudovorticity and vorticity profiles obtained 2.69 s after the beginning of the HVBK simulation. Profiles are averaged either in the vertical (a-c) or horizontal (d-f) directions. Pseudovorticity profiles obtained from (a,d) deuterium D₂ and (b,e) hydrogen H₂ particles. Black, red and blue lines in these panels indicate that corresponding particle radii are 5, 10 and 30 μ m, respectively. (c,f) Pseudovorticity profiles for tracers are plotted as red lines, with black lines indicating corresponding vorticity profiles. Note the different scales on the vertical axes of the right-hand panels for pseudovorticity (left axes) and vorticity (right axes). The reference system is as in figure 11.

(Švančara *et al.* 2020). It then follows that, as mentioned above, the choice we made here for the annulus size is mainly motivated by the requirement of having enough particles in the region of interest for the pseudovorticity calculation.

In order to make a quantitative comparison, we present in figure 12 the root-mean-square profiles of pseudovorticity for the different particle families. These are compared with the corresponding vorticity profiles. We clearly observe that particle inertia becomes more important for the heavier and larger particles, i.e. for the deuterium particles with 30 μ m radii, corresponding to the largest Stokes number. Additionally, vorticity and pseudovorticity profiles qualitatively differ, in agreement with the above discussion.

4.3. Ring position, velocity and radius

Table 2 lists relevant vortex ring properties averaged over the entire HVBK simulation, for times ranging from 0.09 to 26.93 s. The exception is the ring propagation velocity, which is averaged between 1.89 and 25.14 s due to the Gaussian algorithm chosen for its calculation (Švančara *et al.* 2020). Note that the small dependence of the ring velocity on the dataset is most likely related to the above mentioned procedure of ring positioning in the box

Dataset	$\begin{array}{c} \theta_0 \\ (s^{-1}) \end{array}$	$v \pmod{(\text{mm s}^{-1})}$	2 <i>R</i> (mm)	$\frac{C}{(\mathrm{mm}^2 \mathrm{s}^{-1})}$	A_C (mm ²)	St
Vorticity	1.000	5.24 ± 0.77	7.10 ± 0.11	82.86 ± 5.30	11.63 ± 4.22	
Tracer	0.010	5.24 ± 0.78	6.53 ± 0.25	1.51 ± 0.16	35.10 ± 2.96	_
D ₂ 5 μm	0.010	5.24 ± 0.78	6.36 ± 0.21	1.58 ± 0.14	34.70 ± 2.51	0.07
D ₂ 10 μm	0.010	5.24 ± 0.77	5.79 ± 0.17	1.74 ± 0.10	37.37 ± 3.04	0.27
$D_2 30 \mu m$	0.005	5.28 ± 0.84	4.99 ± 0.47	0.74 ± 0.30	51.52 ± 3.34	2.47
$H_2 5 \mu m$	0.010	5.24 ± 0.78	6.63 ± 0.28	1.49 ± 0.16	36.58 ± 3.13	0.04
$H_2 \ 10 \ \mu m$	0.020	5.24 ± 0.77	7.32 ± 0.44	1.24 ± 0.18	25.18 ± 1.79	0.16
$H_2 30 \mu m$	0.050	5.24 ± 0.77	6.34 ± 1.14	1.60 ± 0.37	14.24 ± 2.63	1.45

Table 2. Properties of the HVBK ring. (i) Chosen threshold θ_0 for ring identification, set to approximately 10% of the maximum magnitude at late times. (ii) Ring vertical velocity v; see also figure 13(*d*–*f*). (iii) Ring diameter 2*R*; see also figure 14. (iv) Ring circulation *C* from (3.2). (v) Ring area A_C . (vi) Stokes number $St = \tau_p/\tau_r$, where $\tau_r \approx 17$ ms. Results obtained by applying the chosen threshold values. Symbols as in Švančara *et al.* (2020).

centre at each time step. Indeed, this centring procedure was performed using the normal fluid vorticity maximum, whose magnitude and position do not depend on the particle type.

In figure 13 we present the vortex ring vertical position and velocity, computed from the pseudovorticity maps. For comparison, we also show the results directly obtained from the vorticity field, plotted as black lines. Remarkably, the determination of the ring location is robust and independent of the particle family. Concerning its velocity, the general trend is well reproduced, but some periodic oscillations are visible for the heaviest and larger particles – namely the D_2 30 μ m family, corresponding to the largest Stokes number. These oscillations are likely related to the periodicity of the domain because their period – about 5 s – is approximately the time it takes for the ring to travel through the periodic box. Additionally, as mentioned above, when the ring passes, it leaves in its wake an inhomogeneous distribution of particles that is later encountered in front of the ring. This effect is more marked for particles with high inertia, as is also apparent from figure 10.

The determination of the ring radius, plotted in figure 14, is more sensitive to particle inertia. It is evident from the figure – see especially figure 14(a) – that the ring radius estimate does not depend appreciably on particle size and density for small enough particles, with $St \leq 0.1$. Additionally, for these particles, the expected time increase of the radius, evident from the vorticity trend, is recovered; see also the related discussion in § 3.3.

Another remarkable feature that can be seen in figure 14 is that, for particle radii of 5 and 10 μ m, the ring radius *R* is – on average – larger for the hydrogen particles than for the heavier ones (the ring diameter 2*R* was defined in § 3.3 as the distance between the observed counter-rotating vortices). This may indicate that deuterium particles – heavier than the fluid – tend not to probe the ring core region, as visualized in experiments (figure 1) and in numerical simulations (figure 10). Moreover, for deuterium particles with radii equal to 10 and 30 μ m, the ring radius does not change appreciably with time and is consistently smaller than that observed for tracer particles.

The peculiar behaviour observed in figure 14(b) for the largest hydrogen particles, those with St = 1.45, is at present not entirely understood. However, from the corresponding pseudovorticity maps – not shown here – one can notice that, for sufficiently large times, larger than approximately 10 s, the underlying dipolar ring structure is replaced by two



Figure 13. Temporal evolution of the ring vertical position (a-c) and velocity (d-f) obtained from the HVBK simulation. Panel disposition and line colour code as in figure 12. Note that here, for the sake of clarity and comparison, we plot the absolute values of ring position and velocity in a fixed coordinate system, not moving with the ring and coinciding with the moving frame of reference at the beginning of the simulation (in the coordinate system of figure 11 the ring is moving downward, in the negative direction). The velocity time limits are due to the Gaussian algorithm chosen for the velocity calculation (Švančara *et al.* 2020).

adjacent dipoles. On the other hand, one may more generally say that such large particles are not suitable for tracking the ring.

Note finally that, for the sake of completeness, we also list in table 2 the values of ring circulation and area for the different particle families, in the same fashion as in table 1, but we do not show here the corresponding time trends because they do not add any information on our discussion, that is, they merely confirm the already reported findings.

5. Conclusions

In a recent experimental work (Švančara *et al.* 2020) particle tracking velocimetry was used to construct Lagrangian pseudovorticity maps that were found to provide valuable information on the propagation of macroscopic vortex rings in superfluid ⁴He. In that work, it was also shown analytically that such maps are closely related to the Eulerian vorticity. For such a mathematical proof, it was assumed that particles act as ideal fluid tracers and that they are homogeneously distributed in space. In practice, such assumptions are rarely met in experiments.

In this work, we have numerically studied how particles can be used to study the propagation of vortex rings, in the spirit of Švančara *et al.* (2020). We have performed two sets of numerical simulations. In the first set, we have employed the finite element method to study the behaviour of an isolated vortex ring in a viscous fluid, in a geometry



Figure 14. (a,b) Ring radius as a function of the time *t* elapsed from the beginning of the HVBK simulation. Note the different scales on the vertical axes. Reference system as in figure 13.

similar to that of the experimental setting used by Švančara *et al.* (2020). Fluid tracers were specifically added to the flow to investigate the reliability of pseudovorticity maps. In the second set, we have performed numerical simulations of an isolated vortex ring in superfluid ⁴He, in a periodic domain, using the Hall–Vinen–Bekarevich–Khalatnikov model. In this case, we have considered the dynamics of several particle families of different sizes and densities. We have thus addressed the effect of particle inertia on the propagation properties of an isolated vortex ring.

Overall, our results confirm that the Lagrangian pseudovorticity is a relevant estimator for quantifying general features of isolated vortical structures, such as their position and propagation velocity. This observation is robust and independent of particle inertia. However, we have seen that the intensity of pseudovorticity fields can be much weaker than the actual Eulerian vorticity, especially for sparse particle distributions, and that the ring size is not correctly captured when highly inertial particles are used.

The outcome is especially important in the absence of Eulerian data, which is the case for state-of-the-art visualization experiments in superfluid ⁴He (see e.g. Švančara *et al.* 2021). More generally, our work neatly demonstrates that quantitative comparisons between particle trajectories obtained under different experimental conditions can be performed using pseudovorticity maps, taking also into account particle concentration and inertia. Additionally, future work could focus on establishing the relevance of such a measure for the identification of Lagrangian coherent structures, which is especially relevant for geophysical flows (see e.g. Hadjighasem & Haller 2016).

Acknowledgements. We thank P. Švančara for valuable help. The HVBK computations were carried out on the Mésocentre SIGAMM hosted at the Observatoire de la Côte d'Azur and on the French HPC cluster OCCIGEN through the GENCI allocation A0072A11003.

Funding. J.I.P. and G.K. were supported by the Agence Nationale de la Recherche through the GIANTE ANR-18-CE30-0020-01 project. O.O., M.P., J.H. and M.L.M. acknowledge the support of the Czech Science Foundation (GAČR) under grant no. 19-00939S. M.P. was also supported by the Charles University Research Program no. UNCE/SCI/023.

Declaration of interests. The authors report no conflict of interest.

Author ORCIDs.

- M. Pavelka https://orcid.org/0000-0003-0605-6737;
- ^D M. La Mantia https://orcid.org/0000-0002-7159-5924;
- J.I. Polanco https://orcid.org/0000-0001-7705-753X;
- [®] G. Krstulovic https://orcid.org/0000-0002-9934-6292.

Appendix A. Analytical results

Švančara *et al.* (2020) reported that the Lagrangian pseudovorticity θ is half of the Eulerian vorticity ω . This specifically occurs in the close vicinity of the inspection point *r*, defined in (2.1), and for homogeneous and isotropic distributions of fluid particles, having smooth velocity fields. Although these conditions are at present not met in the case of experiments in He II, the result clearly demonstrates that θ and ω are related, and therefore supports the use of the Lagrangian pseudovorticity, especially in the absence of Eulerian data.

In this appendix we further investigate analytically the relation between pseudovorticity and vorticity in order to explain some results reported in the main text.

Following Švančara et al. (2020) we first rewrite (2.1) as

$$\theta(\mathbf{r}, \mathbf{R}) = \frac{1}{N} \int_{D(\mathbf{r}, \mathbf{R})} f(\mathbf{r}') \frac{\left[\left(\mathbf{r}' - \mathbf{r}\right) \times \mathbf{u}(\mathbf{r}')\right]_z}{|\mathbf{r}' - \mathbf{r}|^2} \,\mathrm{d}^2 \mathbf{r}',\tag{A1}$$

where D(r, R) is a circle, centred at r, with radius R,

$$N = \int_{D(\mathbf{r},R)} f(\mathbf{r}') \,\mathrm{d}^2 \mathbf{r}' \tag{A2}$$

and f denotes the particle distribution function (note that here we disregard the time dependence of the problem).

We now estimate the pseudovorticity for a two-dimensional vortex characterized by a purely azimuthal fluid velocity, having magnitude

$$v_a = \frac{\Gamma}{2\pi r} g(r^2),\tag{A3}$$

where Γ is the constant fluid circulation associated with the vortex and r indicates the distance from the vortex centre. We assume here that $g(r^2)$ is a smooth, non-negative function, vanishing at the origin, $g(r^2) = O(r^2)$ for $r \to 0$, and at most constant at infinity, $g(r^2) = O(1)$ for $r \to \infty$. The corresponding flow vorticity magnitude can then be obtained as

$$\omega = \frac{1}{r} \frac{\partial r v_a}{\partial r} = \frac{\Gamma}{\pi} g'(r^2), \tag{A4}$$

where $g'(r^2)$ indicates the relevant derivative of $g(r^2)$.

The pseudovorticity for a circle of radius R, centred at the inspection point $(x_0, 0)$, in the Cartesian reference system having the origin at the vortex centre, can then be analytically determined, for a given function $g(r^2)$, if also the spatial particle distribution f is known. We first work with an unknown function g and a uniform particle distribution

 $f = N/(\pi R^2)$, N being the number of fluid particles. Using (A3) we can then rewrite (A1) as

$$\theta(x_0; R) = \frac{1}{\pi R^2} \int_{D(x_0, R)} \frac{\mathbf{r} - (x_0, 0)}{|\mathbf{r} - (x_0, 0)|^2} \times \left[\frac{\Gamma}{2\pi r} g(r^2) \mathbf{e}_a\right] \mathrm{d}^2 \mathbf{r},\tag{A5}$$

where e_a indicates the azimuthal velocity direction in the chosen Cartesian coordinate system and r is the magnitude of r.

In the polar reference system having the origin at the inspection point $(x_0, 0)$, (A5) becomes

$$\theta(x_0; R) = \frac{\Gamma}{2\pi^2 R^2} \int_0^R \int_0^{2\pi} \frac{g(r^2)}{r^2} \left(r' + x_0 \cos \varphi'\right) d\varphi' dr',$$
(A6)

where r' and φ' are the chosen polar coordinates and $r^2 = (r')^2 + 2r'x_0 \cos \varphi' + x_0^2$. Since near the origin $g(r^2) = O(r^2) = g'(0)r^2 + O(r^4)$, the integrand in (A6) has no singularity; g'(0) denotes here the derivative of $g(r^2)$ at the vortex centre. Additionally, for $x_0 \to \infty$ the integral in (A6) goes to zero, that is, the pseudovorticity vanishes far from the vortex centre, because the velocity vanishes too.

For $x_0 = 0$, when the inspection point $(x_0, 0)$ coincides with the vortex centre, (A6) becomes

$$\theta(0; R) = \frac{\Gamma}{\pi R^2} \int_0^R \frac{g[(r')^2]}{r'} \,\mathrm{d}r'.$$
 (A7)

Since g behaves at infinity at most as a constant, the integral in (A7) behaves at most as $\ln R$. Therefore, for large R values, $\theta(0; R)$ goes to zero. For small R values, we obtain

$$\lim_{R \to 0} \theta(0; R) = \lim_{R \to 0} \frac{\Gamma}{\pi R^2} \int_0^R \frac{g'(0)(r')^2 + O[(r')^4]}{r'} \, \mathrm{d}r' = \frac{\Gamma}{2\pi} g'(0) = \frac{\omega|_{r=0}}{2}, \quad (A8)$$

which means that, at the origin, the pseudovorticity converges to half of the vorticity. This is compatible with the general result reported by Švančara *et al.* (2020), where it was shown that, for particle distributions radially symmetric around the inspection point, the pseudovorticity converges to half of the vorticity, for sufficiently small R values.

Additionally, using the Hermite–Hadamard inequality (Niculescu & Persson 2006), it can be shown that $\theta(0; R)$ is a decreasing function of R near the origin, because the function $g(r^2)/r$ is there concave. However, for large R values, the latter must become convex, since it is positive, but, at the same time, it is expected that, for large R values, the pseudovorticity goes to zero and therefore its variations are negligible.

In figure 15 we plot $\theta(0; R)$ as a function of *R*, for a Lamb–Oseen vortex, having the azimuthal velocity magnitude

$$v_{LO} = \frac{\Gamma}{2\pi r} g(r^2) = \frac{\Gamma}{2\pi r} \left[1 - \exp\left(\frac{-r^2}{4t\nu}\right) \right],\tag{A9}$$

where r denotes the distance from the vortex centre and, as in figure 2, we set $\Gamma = 50$, t = 10 and v = 0.01 (for the sake of simplicity and generality, we do not explicitly mention in this appendix relevant physical units). We note in passing that the corresponding flow vorticity magnitude can be written as

$$\omega_{LO} = \frac{\Gamma}{4\pi t\nu} \exp\left(\frac{-r^2}{4t\nu}\right). \tag{A10}$$



Figure 15. Pseudovorticity $\theta(0; R)$ as a function of the circle radius R, (A7), for a Lamb–Oseen vortex, (A9), with $\Gamma = 50$, t = 10 and $\nu = 0.01$. Note that the pseudovorticity $\theta(0; R)$ approaches half of the vorticity, $\omega/2 \approx 20$, (A10), for $R \to 0$, vanishes for $R \to \infty$ and is a decreasing function of R. For the sake of simplicity and generality, we do not explicitly mention in this appendix relevant physical units.

In summary, the analytical results reported to date – see especially figure 15 – support the view that the pseudovorticity magnitude becomes smaller if the circular area chosen for its estimate increases, as is also mentioned in the main text.

For $x_0 \neq 0$ the pseudovorticity is given by (A6) and we numerically calculated the integral for the just introduced Lamb–Oseen vortex, that with $\Gamma = 50$, t = 10 and v = 0.01. The integration was carried out using the software Wolfram Mathematica and in figure 16 we plot the obtained result for R = 0.3 (open orange circles). A remarkable agreement with half of the actual flow vorticity (red line) is obtained.

In the same figure the green line, showing a pseudovorticity sign change, corresponds to the green open circles in the main panel of figure 2, that is, it displays the pseudovorticity for the chosen Lamb–Oseen vortex, estimated directly from (2.1), using 1000 equally spaced grid points, up to a radius of 10. Additionally, 100 equally spaced fluid particles were distributed in the chosen annular region, having outer radius equal to 0.3 (the pseudovorticity was not computed in the region centre and was set to zero at the vortex centre). Note also that, as mentioned in § 2, the particles are in this case equally spaced along the radial direction, which means that the chosen particle distribution is not radially symmetric and that the particles are clustered in the vortex centre vicinity.

We can see from (A6) that the integrand is non-negative if

$$r' + x_0 \cos \varphi' \ge 0,\tag{A11}$$

which can be rewritten in the Cartesian coordinate system (x, y) having the origin at the vortex centre as

$$\left(x - \frac{x_0}{2}\right)^2 + y^2 \geqslant \left(\frac{x_0}{2}\right)^2.$$
(A12)

The above equality is satisfied for a circle with radius $x_0/2$, centred at $(x_0/2, 0)$. Points outside this circle contribute positively to the pseudovorticity, while points inside it



Figure 16. Pseudovorticity for a Lamb–Oseen vortex as a function of the distance from the vortex axis (radius). Vortex parameters as in figures 2 and 15, that is, $\Gamma = 50$, t = 10 and $\nu = 0.01$. Black line: vortex azimuthal velocity v_{LO} , (A9); red line: half of the vorticity $\omega_{LO}/2$, (A10); green line: pseudovorticity corresponding to the open green circles in the main panel of figure 2, estimated as described in that figure caption, that is, using directly (2.1); open orange circles: pseudovorticity $\theta(x_0; R)$, (A6), with R = 0.3.

contribute negatively, and the intersection of $D(x_0, R)$ with the positively contributing region is larger than that with the negatively contributing region. Therefore, if the function $g(r^2)/r^2$ behaves nearly as a constant within $D(x_0, R)$, the contribution of the positive region (outside that circle) will be larger than that of the negative region (within that circle), and the pseudovorticity will be positive. However, if the function $g(r^2)/r^2$ is considerably larger in the negative region, the pseudovorticity can become negative. Since the function $g(r^2)/r^2$ is here assumed to be smooth, the pseudovorticity should typically be positive, as indicated in figure 16 (open orange circles).

It then follows that the sign changes of the pseudovorticity displayed in figures 2 and 15 – see also figure 4(b) – are most likely an effect of the discrete particle distributions employed for the θ estimates based on (2.1), especially if one also considers that the analytical results reported here are only valid for uniform particle distributions, that is, we set $f = N/(\pi R^2)$ in (A1). Indeed, we would like to address analytically the effect of particle distribution non-uniformities in future studies.

Another possible line of future research could be based on the following result, which can be considered a generalization of the analytical relation between θ and ω reported by Švančara *et al.* (2020). The latter was obtained by expanding the fluid velocity to the second order with respect to the difference (r' - r). If, in a similar fashion, we expand it to the fourth order, we get

$$\theta(\mathbf{r}; \mathbf{R}) = \frac{\omega(\mathbf{r})}{2} + \frac{R^2}{32} \Delta \omega(\mathbf{r})$$
(A13)

for a smooth velocity field, in the close vicinity of the inspection point r, for radially symmetric particle distributions. The pseudovorticity is thus approximately a quadratic function of R and its behaviour depends on the Laplacian of the vorticity at the inspection point. It then follows that, if $\theta(r; R)$ is known, one could estimate the flow vorticity

 $\omega(r)$ using (A13). The latter estimate could be performed by numerical means, or even analytically, and this could be the focus of future studies, assessing in more detail the effect of *R* on the relation between θ and ω .

REFERENCES

- ALNÆS, M.S., BLECHTA, J., HAKE, J., JOHANSSON, A., KEHLET, B., LOGG, A., RICHARDSON, C., RING, J., ROGNES, M.E. & WELLS, G.N. 2015 The FEniCS project version 1.5. Arch. Num. Soft. 3, 9–23.
- BALACHANDAR, S. & EATON, J.K. 2010 Turbulent dispersed multiphase flow. Annu. Rev. Fluid Mech. 42, 111–133.
- BARENGHI, C.F., SKRBEK, L. & SREENIVASAN, K.R. 2014 Introduction to quantum turbulence. *Proc. Natl Acad. Sci. USA* 111, 4647–4652.
- BORNER, H. & SCHMIDT, D.W. 1985 Investigation of large-scale vortex rings in He II by acoustic measurements of circulation. In *Flow of Real Fluids* (ed. G.E.A. Meier & F. Obermeier), *Lecture Notes in Physics*, vol. 235, pp. 135–146. Springer.
- BREZZI, F. & FORTIN, M. 1991 Mixed and Hybrid Finite Element Methods. Springer.
- DIDDEN, N. 1979 On the formation of vortex rings: rolling-up and production of circulation. Z. Angew. Math. Phys. 30, 101–116.
- DONNELLY, R.J. 2009 The two-fluid theory and second sound in liquid helium. Phys. Today 62 (10), 34-39.
- DONNELLY, R.J. & BARENGHI, C.F. 1998 The observed properties of liquid helium at the saturated vapor pressure. J. Phys. Chem. Ref. Data 27, 1217–1274.
- DUDA, D., LA MANTIA, M. & SKRBEK, L. 2017 Streaming flow due to a quartz tuning fork oscillating in normal and superfluid ⁴He. *Phys. Rev.* B **96**, 024519.
- DUDA, D., ŠVANČARA, P., LA MANTIA, M., ROTTER, M. & SKRBEK, L. 2015 Visualization of viscous and quantum flows of liquid ⁴He due to an oscillating cylinder of rectangular cross section. *Phys. Rev.* B **92**, 064519.
- GAN, L. & NICKELS, T.B. 2010 An experimental study of turbulent vortex rings during their early development. J. Fluid Mech. 649, 467–496.
- GHARIB, M., RAMBOD, E. & SHARIFF, K. 1998 A universal time scale for vortex ring formation. J. Fluid Mech. 360, 121–140.
- GLEZER, A. 1988 On the formation of vortex rings. Phys. Fluids 31, 3532-3542.
- GUERMOND, J.L., MINEV, P. & SHEN, J. 2006 An overview of projection methods for incompressible flows. Comput. Meth. Appl. Mech. Engng 195, 6011–6045.
- HADJIGHASEM, A. & HALLER, G. 2016 Level set formulation of two-dimensional Lagrangian vortex detection methods. *Chaos* 26, 103102.
- VAN HINSBERG, M.A.T., THIJE BOONKKAMP, J.H.M., TOSCHI, F. & CLERCX, H.J.H. 2012 On the efficiency and accuracy of interpolation methods for spectral codes. *SIAM J. Sci. Comput.* 34, B479–B498.
- HOMANN, H., KAMPS, O., FRIEDRICH, R. & GRAUER, R. 2009 Bridging from Eulerian to Lagrangian statistics in 3D hydro- and magnetohydrodynamic turbulent flows. *New J. Phys.* **11**, 073020.
- LA MANTIA, M. & SKRBEK, L. 2014 Quantum, or classical turbulence? Europhys. Lett. 105, 46002.
- LANDAU, L. 1941 Theory of the superfluidity of helium II. Phys. Rev. 60, 356-358.
- L'VOV, V.S., NAZARENKO, S.V. & SKRBEK, L. 2006 Energy spectra of developed turbulence in helium superfluids. J. Low Temp. Phys. 145, 125–142.
- MAXEY, M.R. 1987 The gravitational settling of aerosol particles in homogeneous turbulence and random flow fields. *J. Fluid Mech.* **174**, 441–465.
- MAXWORTHY, T. 1974 Turbulent vortex rings. J. Fluid Mech. 64, 227–239.
- MONGIOVÌ, M.S., JOU, D. & SCIACCA, M. 2018 Non-equilibrium thermodynamics, heat transport and thermal waves in laminar and turbulent superfluid helium. *Phys. Rep.* **726**, 1–71.
- NICULESCU, C.P. & PERSSON, L.-E. 2006 Convex Functions and their Applications. Springer.
- NOBACH, H. & BODENSCHATZ, E. 2009 Limitations of accuracy in PIV due to individual variations of particle image intensities. *Exp. Fluids* **47**, 27–38.
- OUTRATA, O. 2020 Numerical methods for vortex dynamics. Master's thesis, Charles University, Prague, Czech Republic.
- POLANCO, J.I. & KRSTULOVIC, G. 2020 Inhomogeneous distribution of particles in coflow and counterflow quantum turbulence. *Phys. Rev. Fluids* **5**, 032601.
- RAFFEL, M., WILLERT, C.E., SCARANO, F., KÄHLER, C.J., WERELEY, S.T. & KOMPENHANS, J. 2018 Particle Image Velocimetry. Springer.

- SERGEEV, Y.A. & BARENGHI, C.F. 2009 Particles-vortex interactions and flow visualization in ⁴He. J. Low Temp. Phys. 157, 429–475.
- SULLIVAN, I.S., NIEMELA, J.J., HERSHBERGER, R.E., BOLSTER, D. & DONNELLY, R.J. 2008 Dynamics of thin vortex rings. J. Fluid Mech. 609, 319–347.
- ŠVANČARA, P., et al. 2021 Ubiquity of particle-vortex interactions in turbulent counterflow of superfluid helium. J. Fluid Mech. 911, A8.
- ŠVANČARA, P., HRUBCOVÁ, P., ROTTER, M. & LA MANTIA, M. 2018 Visualization study of thermal counterflow of superfluid helium in the proximity of the heat source by using solid deuterium hydride particles. *Phys. Rev. Fluids* **3**, 114701.
- ŠVANČARA, P. & LA MANTIA, M. 2017 Flows of liquid ⁴He due to oscillating grids. J. Fluid Mech. 832, 578–599.

ŠVANČARA, P. & LA MANTIA, M. 2019 Flight-crash events in superfluid turbulence. J. Fluid Mech. 876, R2.

- ŠVANČARA, P., PAVELKA, M. & LA MANTIA, M. 2020 An experimental study of turbulent vortex rings in superfluid ⁴He. J. Fluid Mech. 889, A24.
- VILLOIS, A., PROMENT, D. & KRSTULOVIC, G. 2020 Irreversible dynamics of vortex reconnections in quantum fluids. *Phys. Rev. Lett.* **125**, 164501.
- VINEN, W.F. & NIEMELA, J.J. 2002 Quantum turbulence. J. Low Temp. Phys. 128, 167-231.
- WACKS, D.H., BAGGALEY, A.W. & BARENGHI, C.F. 2014 Large-scale superfluid vortex rings at nonzero temperatures. *Phys. Rev.* B **90**, 224514.